

Patient Information Leaflet

Glandomed with Lidocaine 0.1% (=100 mg Lidocaine HCl in 100 ml Glandomed)

This oral mouth rinse can be prepared for you by a pharmacist.
Please read all of this leaflet carefully before you start taking this medicine.
If you have any further questions ask your doctor or pharmacist.

For oral rinse, not for drinking

Composition:

500 ml solution contains:

Medicinally active ingredient

0.5 g lidocaine hydrochloride

Other ingredients:

Macrogol, sodium hydrogen carbonate, sodium edetate, chlorhexidine digluconate (for preservation), orange flavouring.

Dosage form and content:

Bottle with 500 ml solution for rinsing the oral mucosa

Substance or indication group:

Topical anaesthetic

Fields of application:

For anaesthesia of the oral mucosa in incipient and advanced mucositis (inflammation of the oral mucosa).

Contraindications:

Glandomed with lidocaine must not be used in:

Intolerance to lidocaine

Severe disorders of the stimulus-forming and stimulus-conducting system in the heart

The contraindications do not apply to Glandomed without lidocaine.

Interactions:

There is no information available

Dosage instructions, method and duration of use:

Rinse the mouth with 10 to 15 ml of the indiluted solution for 2 minutes, then spit out, do not rinse the mouth with water after, do not eat or drink for 15 minutes. The application can be done hourly if necessary

Side effects:

Contact allergy in case of hypersensitivity to lidocaine

Notes:

- Use only clear solution from intact container
- Keep medicines out of reach of children.
- the numbness on the mucous membrane is a natural reaction to lidocaine, it is reversible
- Do not drink or swallow.
- Shelf life: 4 weeks from production, discard leftovers
- prescription medicine

Caution:

The bottle must be provided with a child-resistant closure because accidentally drinking the entire content can have serious to fatal consequences.

For preparation:

www.mucoisitis.de/recipes

Version 2021/03